# Supporting Information 2 for:

# Automated Purification of DNA Origami with SPRI Beads

Chalmers Chau<sup>1,2</sup> \*<sub>‡</sub>, Gayathri Mohanan<sup>1,2</sup><sub>‡</sub>, Iain Macaulay<sup>3,4</sup>, Paolo Actis<sup>1,2</sup>, Christoph Wälti<sup>1,2</sup> \*

<sup>1</sup> School of Electronic and Electrical Engineering, University of Leeds, LS2 9JT, UK

<sup>2</sup> Bragg Centre for Materials Research, University of Leeds, LS2 9JT, UK

<sup>3</sup> Earlham Institute, Norwich Research Park, Norwich, NR1 7UZ, UK

<sup>4</sup> School of Biological Sciences, University of East Anglia, Norwich, Norfolk, NR4 7TJ, UK

<sup>‡</sup> These authors contribute equally.

\* Corresponding: <u>c.c.chau@leeds.ac.uk;</u> <u>c.walti@leeds.ac.uk;</u>

### Intact origami counting

Throughout this section, for origamis that are counted as intact, a black number will be used. For origamis that are counted as damaged, a red number will be used.



Below shows an AFM image as an example:

In this example, a total of 17 intact origamis and 4 damaged origamis.

### S-400 HR

Images below were used to quantify the number of intact origamis purified using S-400 HR spin column.





### SPRI - 0.8X ratio

Images below were used to quantify the number of intact origamis purified using SPRI at 0.8X ratio column.







### **MWCO – 1**

Images below were used to quantify the number of intact origamis purified using 100 kDa MWCO – 1 concentrator.







### **MWCO – 2**

Images below were used to quantify the number of intact origamis purified using 100 kDa MWCO – 2 concentrator.



# **Ethanol Precipitation**

Images below were used to quantify the number of intact origamis purified using ethanol precipitation.









## **PEG precipitation**

Images below were used to quantify the number of intact origamis purified using PEG precipitation.











### **Phase separation**

Images below were used to quantify the number of intact origamis purified using phase separation method.





### Size Exclusion Chromatography

Images below were used to quantify the number of intact origamis purified using size exclusion chromatography.







Height Sensor

200.0 nm





2.7<u>n</u>m

-1.6 nm

Height Sensor

200.0 nm



Height Sensor

200.0 nm

### Intact CRP functionalised DNA origami count

Throughout this document, for origamis that are counted as intact, a green circle will be used. For origamis that are counted as damaged, a red circle will be used. For origamis that are intact but no CRP, a blue circle will be used.

Below shows an AFM image as an example:

# Green – CRP bound origami – 11 Blue – intact but no CRP - 9 Red – deformed (not considered in counting) - 10



In this example, a total of 11 intact origamis with CRP, 9 intact origamis and 10 damaged origamis.

### **SPRI cleaned DNA origamis**

Green – CRP bound origami – 11 Blue – intact but no CRP - 9 Red – deformed (not considered in counting) - 10



220.0 nm

Green – CRP bound origami – 12 Blue – intact but no CRP - 4 Red – deformed (not considered in counting) - 6



220.0 nm

Green – CRP bound origami – 6 Blue – intact but no CRP - 22 Red – deformed (not considered in counting) - 16



210.0 nm

Green – CRP bound origami – 4 Blue – intact but no CRP - 7 Red – deformed (not considered in counting) – 1



210.0 nm

Green – CRP bound origami – 3 Blue – intact but no CRP - 7 Red – deformed (not considered in counting) – 5



210.0 nm

Green – CRP bound origami – 9 Blue – intact but no CRP - 3 Red – deformed (not considered in counting) – 1



210.0 nm